

## Exhibit 2-8: Non-Drug Treatment Options for Prostate Cancer

<b>Treatment</b>	<b>Overview</b>	<b>Risks</b>	<b>Recovery</b>	<b>Efficacy</b>	<b>Cost</b>
Radical Prostatectomy (RP)	Surgical procedure to remove the entire prostate gland and surrounding tissue	Urinary tract obstruction; incontinence; impotence	3-5 Weeks	78% negative long-term biopsy	\$25K-\$30K
Targeted Cryosurgery	MIS procedure that freezes prostate gland to destroy cancerous cells	Incontinence; impotence; urinary tract obstruction	< 1 Week	82% negative long-term biopsy	\$12-15K
External Beam Radiation	1-2 month treatment regimen to stop growth of cancerous cells	Abnormal bowel function; urinary tract obstruction; impotence; incontinence	Fatigue 1-2 months following last treatment	56% negative long-term biopsy	\$12-15K
Brachytherapy	Permanent implantation of approximately 100 radioactive seeds into the prostate gland	Radiation poisoning; irritated urinary function; impotence	Pain-free after 1 week	78% negative long-term biopsy	\$12-15K

Source: MedMarket Diligence, LLC

Report #D101: The U.S. Therapeutic Oncology Market, 2003-2013: Trends, Technologies & Opportunities (See <http://www.mediligence.com/rpt-d101.htm>)

For more details, contact: Patrick J. Driscoll, <mailto:patrick@mediligence.com>  
 MedMarket Diligence, LLC  
 Tel: 949.859.3401